

Selection for Fitness vs. Selection for Robustness in RNA Secondary Structure Folding

Claus O. Wilke
Digital Life Laboratory
California Institute of Technology, Mail-Code 136-93
Pasadena, CA 91125
wilke@caltech.edu

February 2, 2008

Abstract

We investigate the competition between two quasispecies residing on two disparate neutral networks. Under the assumption that the two neutral networks have different topologies and fitness levels, it is the mutation rate that determines which quasispecies will eventually be driven to extinction. For small mutation rates, we find that the quasispecies residing on the neutral network with the lower replication rate will disappear. For higher mutation rates, however, the faster replicating sequences may be outcompeted by the slower replicating ones in case the connection density on the second neutral network is sufficiently high. Our analytical results are in excellent agreement with flow-reactor simulations of replicating RNA sequences.

Keywords: quasispecies, mutant cloud, neutral networks, RNA secondary structure folding, selection of robustness

At high mutation rates, the number of mutated offspring generated in a population far exceeds the number of offspring identical to their parents. As a result, a stable cloud of mutants, a so-called quasispecies (Eigen and Schuster 1979; Eigen et al. 1988; Eigen et al. 1989; Nowak 1992; Wilke et al. 2001), forms around the fastest replicating genotypes. Experimental evidence in favor of such a persistent cloud of mutants is available from RNA viruses (Steinhauer et al. 1989; Domingo and Holland 1997; Burch and Chao 2000) and in vitro RNA

replication (Biebricher 1987; Biebricher and Gardiner 1997); both are cases in which a high substitution rate per nucleotide is common (Drake 1993). The existence of a quasispecies has important implications for the way in which selection acts, because the evolutionary success of individual sequences depends on the overall growth rate of the quasispecies they belong to. As a consequence, organisms with a high replication rate that produce a large number of offspring with poor fitness can be outcompeted by organisms of smaller fitness that produce a larger number of also-fit offspring (Schuster and Swetina 1988). Similarly, if a percentage of the possible mutations is neutral, and the majority of the non-neutral mutations is strongly deleterious, then the growth rate of a quasispecies depends significantly on the connection density (the number of nearby neutral mutants of an average viable genotype) of the neutral genotypes (van Nimwegen et al. 1999). Therefore, a neutral network (a set of closely related mutants with identical fitness) with high connectivity can be advantageous over one with higher fitness, but lower connectivity. Here, we are interested in this latter possibility. In particular, we investigate the competition of two quasispecies residing on separate neutral networks with different connection densities and replication rates, and determine under what conditions selection favors the more fit (i.e., of higher replication rate) or the more robust (more densely connected) mutant cloud. Our approach is closely related to the study of holey landscapes, in which all genotypes are classified into either viable or inviable ones (Gavrilets and Gravner 1997; Gavrilets 1997).

However, we extend this picture by further subdividing the viable genotypes into two groups with different replication rates.

The paper is organized as follows. First, we describe a simple model of a quasispecies on a single neutral network, and demonstrate that the model is consistent with simulations of RNA sequences. Then, based on this model, we present a model of two competing quasispecies, and compare the second model with simulation results as well. Following that, we study the probability of fixation of a single advantageous mutant that arises in a fully formed quasispecies. Finally, we discuss the implications of our results and give conclusions.

POPULATION DYNAMICS ON A SINGLE NEUTRAL NETWORK

Before we can address the competition of two quasispecies, we need a good description of a single quasispecies on a neutral network. A fundamental contribution to this problem has been made by van Nimwegen et al. (1999), who showed that the average fitness of a population on a neutral network is determined only by the fitness of the neutral genotypes, the mutation rate, and the largest eigenvalue of the neutral genotypes' connection matrix. The connection matrix is a symmetric matrix with one row/column per neutral genotype. It holds a one in those positions where the row- and the column-genotype are exactly one point-mutation apart, and a zero otherwise. In theory, the formalism of van Nimwegen et al. describes a population on a neutral network well. However, the exact connection matrix is normally not known, which implies that we cannot calculate the population dynamics from first principles. Nevertheless, we can base a very simple model on the fact – also established by van Nimwegen et al. – that the average neutrality in the population, which is exactly the largest eigenvalue of the connection matrix, is independent of the mutation rate. The main assumption of our simple model is that the population behaves as if all sequences in the population had the same neutrality ν , where ν is given by the average neutrality in the population. Moreover, we consider genetic sequences of length l , and assume a per-symbol copy fidelity of q . Then, the effective copy fidelity or neutral fidelity (Ofria and Adami 2001) Q , i.e., the probability with which on average a viable

sequence gives birth to offspring that also resides on the neutral network, is given by

$$Q = [1 - (1 - q)(1 - \nu)]^l \approx e^{-l(1-q)(1-\nu)}. \quad (1)$$

Now, we can devise a two-concentration model in which $x_1(t)$ is the total concentration of all sequences on the neutral network, and $x_d(t)$ is the concentration of sequences off the network (these sequences are assumed to replicate so slowly that their offspring can be neglected). The two quantities satisfy the equations

$$\begin{aligned} \dot{x}_1(t) &= w_1 Q x_1(t) - e(t) x_1(t), \\ \dot{x}_d(t) &= w_1 (1 - Q) x_1(t) - e(t) x_d(t), \end{aligned} \quad (2a)$$

where w_1 is the fitness of the sequences on the neutral network, and $e(t)$ is the excess production (or mean fitness in the population) $e(t) = w_1 x_1(t)$. Equation (2a) can be integrated directly. We find

$$x_1(t) = \frac{Q x_1(0)}{x_1(0) + [Q - x_1(0)] e^{-w_1 Q t}}. \quad (3)$$

In the steady state ($t \rightarrow \infty$), this implies that the concentration of sequences on the network is equal to the effective fidelity Q ,

$$x_1 = Q = e^{-l(1-q)(1-\nu)}. \quad (4)$$

Therefore, by measuring the decay of the concentration of sequences on the neutral network as a function of the copy fidelity q , we can estimate the population neutrality ν .

Note that the above description of the evolving population is similar to the one presented by Reidys et al. (2001), with one important conceptual difference. The article by Reidys et al. (2001) was completed before van Nimwegen et al.'s work was available, and therefore it was not clear what their effective fidelity did actually relate to. Here, on the other hand, we know that Q depends only on the copy fidelity per nucleotide, q , and the average population neutrality ν , which is independent of q and could be calculated exactly if the connection matrix of the neutral genotypes was known.

We have measured the average equilibrium concentration x_1 of sequences on the network for RNA

secondary structure folding. RNA folding is a reliable test case, and has been applied to a wide array of different questions related to the dynamics of evolution (Fontana et al. 1993; Huynen et al. 1996; Fontana and Schuster 1998; Schuster and Fontana 1999; Ancel and Fontana 2000; Reidys et al. 2001). We simulated a flow reactor using the Gillespie algorithm (Gillespie 1976), and performed the RNA folding with the Vienna package (Hofacker et al. 1994), version 1.3.1, which uses the parameters given by Walter et al. (1994). The carrying capacity was set to $N = 1000$ sequences, and the reactor was initially filled with 1000 identical copies of a sequence that folded into a given target structure. Sequences folding into the target structure were replicating with rate one per unit time, and all other sequences with rate 10^{-6} per unit time. We let the reactor equilibrate for 50 time steps, and then measured the average concentration of correctly folding sequences over the next 150 time steps.

Results for the two different target structures depicted in Fig. 1 are shown in Fig. 2. In both cases, we see a very clear exponential decay. Up to a mutation rate of 0.05, which is quite high for the sequences of length $l = 62$ we are considering here, we cannot make out a significant deviation from a straight line in the log-linear plot. This verifies the applicability of our simple model to evolving RNA sequences. Note that our simulations also show a significant difference in the effective neutrality of the two structures, which will be of importance in the next section.

TWO COMPETING QUASISPECIES

Analytical Model

Above, we have established a simple description for a quasispecies residing on a single neutral network. In a similar fashion, we can treat the competition of two quasispecies residing on separate networks. We classify all sequences into three different groups: sequences on network one, sequences on network two, and dead sequences (sequences that replicate much slower than sequences on either of the two networks, or do not replicate at all). We denote the respective relative concentrations by x_1 , x_2 , and x_d . We make the further assumption that all sequences within a neutral network i have the same probability Q_i to mutate into another sequence on network

i , and we neglect mutations from one network to the other. The probability to fall off of a network i is hence $1 - Q_i$. The differential equations for an infinite population are then:

$$\dot{x}_1(t) = w_1 Q_1 x_1(t) - e(t) x_1(t), \quad (5a)$$

$$\dot{x}_2(t) = w_2 Q_2 x_2(t) - e(t) x_2(t), \quad (5b)$$

$$\dot{x}_d(t) = w_1(1 - Q_1)x_1(t) + w_2(1 - Q_2)x_2(t) - e(t)x_d(t), \quad (5c)$$

where w_1 and w_2 are the fitnesses of sequences on network one or two, respectively, and $e(t)$ is the excess production $e(t) = w_1 x_1(t) + w_2 x_2(t)$. In order to solve Eq. (5), it is useful to introduce the matrix

$$\mathbf{W} = \begin{pmatrix} w_1 Q_1 & 0 & 0 \\ 0 & w_2 Q_2 & 0 \\ w_1(1 - Q_1) & w_2(1 - Q_2) & 0 \end{pmatrix}. \quad (6)$$

We further need the exponential of $\mathbf{W}$, which is given by

$$\exp(\mathbf{W}t) = \begin{pmatrix} e^{w_1 Q_1 t} & 0 & 0 \\ 0 & e^{w_2 Q_2 t} & 0 \\ \frac{1-Q_1}{Q_1}(e^{w_1 Q_1 t} - 1) & \frac{1-Q_2}{Q_2}(e^{w_2 Q_2 t} - 1) & 1 \end{pmatrix} \quad (7)$$

Now, if we combine the concentrations x_1 , x_2 , x_d into a vector $\mathbf{x} = (x_1, x_2, x_d)^t$, we find

$$\mathbf{x}(t) = \exp(\mathbf{W}t) \cdot \mathbf{x}(t) / [\hat{\mathbf{e}} \cdot \exp(\mathbf{W}t) \cdot \mathbf{x}(0)] \quad (8)$$

with $\hat{\mathbf{e}} := (1, 1, 1)$. The denominator on the right-hand side of Eq. (8) corresponds to the cumulative excess production $e_{\text{cum}}(t) = \int_0^t e(t) dt$, which is given by

$$\begin{aligned} e_{\text{cum}}(t) &= \hat{\mathbf{e}} \cdot \exp(\mathbf{W}t) \cdot \mathbf{x}(0) \\ &= \frac{x_1(0)}{Q_1}(e^{w_1 Q_1 t} + Q_1 - 1) \\ &\quad + \frac{x_2(0)}{Q_2}(e^{w_2 Q_2 t} + Q_2 - 1) + x_d(0). \end{aligned} \quad (9)$$

The solution to Eq. (5) follows now as

$$x_1(t) = \frac{e^{w_1 Q_1 t}}{e_{\text{cum}}(t)} x_1(0), \quad (10a)$$

$$x_2(t) = \frac{e^{w_2 Q_2 t}}{e_{\text{cum}}(t)} x_2(0), \quad (10b)$$

$$\begin{aligned} x_d(t) &= \frac{1}{Q_1 Q_2 e_{\text{cum}}(t)} \left[(e^{w_1 Q_1 t} - 1)(1 - Q_1) Q_2 x_1(0) \right. \\ &\quad \left. + (e^{w_2 Q_2 t} - 1)(1 - Q_2) Q_1 x_2(0) + Q_1 Q_2 x_d(0) \right]. \end{aligned} \quad (10c)$$

There exist two possible steady states. If $w_1 Q_1 > w_2 Q_2$, then for $t \rightarrow \infty$ we have $x_1 = Q_1$, $x_2 = 0$, $x_d = 1 - Q_1$. If $w_1 Q_1 < w_2 Q_2$, on the other hand, the steady state distribution is given by $x_1 = 0$, $x_2 = Q_2$, $x_d = 1 - Q_2$. The most interesting situation occurs when for a given w_1 and w_2 , the steady state depends on the mutation rate. This happens if $w_1 > w_2$, but $\nu_1 < \nu_2$, or vice versa. Namely, if we express Q_i as given in Eq. (1), we obtain from $w_1 Q_1 = w_2 Q_2$ the critical copy fidelity

$$q_c = 1 - \frac{\ln(w_2/w_1)}{l(\nu_1 - \nu_2)}. \quad (11)$$

Clearly, q_c can only be smaller than one if either $w_1 > w_2$ and $\nu_1 < \nu_2$ or vice versa. Therefore, this is a necessary (though not sufficient) condition for the existence of two qualitatively different steady states in different mutational regimes. In the language of physics, the transition from one of the two steady state to the other is a first order phase transition (Stanley 1971). The transition is of first order because the order parameter (which we can define to be either x_1 or x_2) undergoes a discontinuous jump from a finite value to zero at the critical mutation rate.

The two phases are not just a mathematical curiosity, they have important biological interpretations. The phase in which the sequences with the larger w_i survive can be considered the “normal” selection regime, i.e., selection which favors faster replicating individuals. We will refer to this situation as the phase of “selection for replication speed”. In the other phase, however, the situation is exactly reversed, and the sequences with the lower intrinsic replication rate w prevail. In this phase, the amount of neutrality (or the robustness against mutations) is more important, and we will consequently refer to this situation as the phase of “selection for robustness”. In Fig. 3, we show two example phase diagrams. These diagrams demonstrate that the selection for robustness is not a pathological situation occurring only for extremely rare sets of parameters, but that in fact both phases have to be considered on equal grounds, none of them can be singled out as the more common one. In particular, as the ratio between w_1 and w_2 approaches unity, the selection for robustness becomes more and more important.

Simulation Results

As in the case of a single quasispecies on a neutral network, we have tested our predictions with simulations of self-replicating RNA sequences in a flow reactor. We assumed that sequences folding into either Fold 1 or 2 (Fig. 1) were replicating with rates $w_1 = 1$ and $w_2 = 1.1$, respectively, while all other folds had a vanishing replication rate. In all results presented below, we initialized the flow reactor with 50% of the sequences folding into Fold 1, and the remaining sequences folding into Fold 2.

Figure 4 shows a comparison between Eq. (10) and four example runs. Apart from finite size fluctuations, which are to be expected in a simulation with $N = 1000$, the analytic expression predicts the actual population dynamics well.

In Fig. 5, we present measurements of the concentrations $x_1(t)$ and $x_2(t)$ as functions of the mutation rate $1 - q$, for a fixed time $t = 200$. The points represent results averaged over 25 independent simulations, and the lines stem from Eq. (10). In agreement with the predictions from our model, we observe two selection regimes, one in which the faster replicating sequences dominate, and one in which the sequences with the higher neutrality have a selective advantage. The transition between the two phases occurs in this particular case approximately at $q = 0.98$, and both the analytical model and the simulations agree well on this value. As is typical for a phase transition, the fluctuations close to the transition point increase significantly, and the time until either of the two quasispecies has gone extinct diverges (the latter point can be seen from the fact that close to the transition point, the disadvantageous fold is still present in a sizeable amount, while further away it has already vanished completely from the population).

Figure 5 also shows that for very small populations, the predictive value of the differential equation approach diminishes, presumably because the choice of a single effective copy fidelity Q is not justified anymore once a minimum population size has been reached. However, as long as we are dealing with population sizes of several hundreds or more, our analytical calculations predict the simulation results very well.

Probability of Fixation

In the previous subsection, we have established that selection acts on the product of replication rate

w and fidelity Q , rather than on the replication rate alone. In particular, for an appropriate choice of parameters, sequences with a lower replication rate can outcompete those with a higher replication rate. However, the competition experiments that we conducted in the previous section were unrealistic in so far that we assumed equal initial concentrations of the two competing types of sequences. A more realistic assumption is that one type (the one with the lower product $w_i Q_i$) dominates the population, while the second type is initially represented through only a single individual. The idea behind this scenario is of course that the second type (with higher product $w_i Q_i$) has arisen through a rare mutation. The question in this context is whether the second type will be able to dominate the population, i.e., whether it will become fixated.

In a standard population genetics scenario, the answer to the above question is simple. If two sequences replicate with w_1 and w_2 , respectively, and mutations between the two sequences can be neglected, then a single sequence of type 2 ($w_2 > w_1$) will become fixated in a background of sequences of type 1 with probability $\pi = 1 - e^{-2s} \approx 2s$, where $s = w_2/w_1 - 1$ is the selective advantage of the newly introduced sequence type (Haldane 1927; Kimura 1964; Ewens 1979). Note, however, that this celebrated result is only correct for a generational model with discrete time steps. In a continuous time model, the equivalent result reads $\pi = s/(1 + s)$. This formula follows from the solution to the problem of the Gambler's Ruin (Feller 1968; Lenski and Levin 1985) when taking the limit of a large population size.

Here, we are not dealing with individual sequences replicating with rate w_i , but rather with quasispecies that grow with rate $w_i Q_i$. A naive way to calculate the fixation probability in this case is simply to replace w_i with $w_i Q_i$ in the expression for the selective advantage, and hope that the result is correct. However, it is not clear from the outset that this approach will work, because the factor Q_i depends on the assumption that a fully developed quasispecies with the appropriate mean neutrality is already present. A single sequence struggling for fixation does not satisfy this condition. Therefore, the actual fixation probability might deviate from the one thus calculated, in particular in circumstances in which a sequence with smaller replication rate is supposed to overtake

an established quasispecies of sequences with higher replication rate.

We performed fixation experiments in both the “selection for replication speed” and the “selection for robustness” phase, in order to clarify whether the naive approach works. In both phases, we allowed a population of size $N = 1000$ to equilibrate, and then introduced a single sequence of the supposedly advantageous type. After 500 time steps, we determined whether the advantageous type had vanished from the population or grown to a significant proportion. By repeating this procedure 100 times, we obtained an estimate for the probability of fixation. As in the previous section, we used $w_1 = 1$ and $w_2 = 1.1$.

In Fig. 6, we compare our simulation results to the predicted fixation probability $\pi = s/(1 + s)$. Within the accuracy of our results, both agree well. This is particularly interesting for mutation rates above 0.02, where we introduce a sequence of lower replication rate into a background of faster replicating sequences. The increased neutrality of the introduced sequence is sufficient to let it rise to fixation in a significant proportion of cases. Moreover, the product $w_i Q_i$ is the sole determinant of the fixation probability. Whether the value of the product $w_i Q_i$ comes mainly from the intrinsic growth rate w_i of the sequences or from the effective fidelity Q_i does not have an observable influence on the dynamics.

DISCUSSION

The good agreement between our analytical model and our simulation results demonstrates that RNA sequences evolving on a neutral network of identical secondary structure folds are well described by only two parameters, their intrinsic replication rate w and their effective copy fidelity Q . In the particular context of two competing distinct folds, we find furthermore that only the product of w and Q is of importance. Indeed, it follows from Eq. (10) that the ratio between $x_1(t)$ and $x_2(t)$ depends only on the respective products of w and Q , but not on the individual values themselves.

Unlike the intrinsic replication rate w , which is a property of the individual, the effective fidelity Q is a group property, as it is given by the average over all sequences in the population of the probability not to “fall off” the neutral network. Thus, in the regime in which Q dominates the evolutionary dynamics (the

phase of selection for robustness in Fig. 3), the evolutionary success of an individual sequence depends strongly on the properties of the group it belongs to. In other words, we find that selection acts on the whole group of mutants, rather than on individuals, despite the absence of standard factors supporting group selection such as spatial subdivision of the population (Wilson 1979), altruistic behavior, parental care (Maynard Smith 1993), or mutual catalytic support (Alves et al. 2001). Here, a sequence with a comparatively high neutrality embedded into a neutral network with a poor overall connection density will be at a disadvantage with respect to a sequence with a comparatively low neutrality that is, however, part of a neutral network with high connection density. The overall higher fidelity of a population on the second network results in a larger fraction of sequences that actually reside on the network, which in turn increases the chance that a particular sequence will be generated as mutant offspring from some other sequence. Moya et al. (2000) noted that this type of group selection should follow from the quasispecies equations, and that populations under this type of selection would be best described by an effective group replication rate r . In the present work, we have shown that this is indeed the case, and we can also derive r (which is simply $r = wQ$) from the quasispecies equations. Namely, the fact that the population neutrality ν (which determines Q) is given by the largest eigenvalue of the connection matrix of neutral genotypes is a direct consequence of the quasispecies equations (van Nimwegen et al. 1999).

Schuster and Swetina (1988) were the first to point out that at high mutation rates, the quasispecies around the highest peak in the landscape can disappear. They focused on situations in which the highest and the second-highest peak in a landscape were of almost equal height, while the immediate mutational neighborhood of the second peak was less deleterious than the one of the first peak. As a consequence, their results seemed to imply that the phase of 'selection of robustness' was only important in the case of very similar peaks. Our results, on the other hand, show that the difference in peak height can be dramatic, if balanced by an equally dramatic difference in robustness.

While our analytical results apply strictly speaking only to infinite populations, we have seen that in sim-

ulations for population sizes as small as $N = 500$, the differential equation approach works well. Moreover, in our experiments on the probability of fixation, we have seen that even very small numbers of the advantageous group (in the extreme only a single sequence) can rise to fixation, despite their intrinsic replication rate being smaller than that of the currently dominating group. This result seems somewhat unintuitive at first, but can be easily understood. The most important aspect of every fixation event is the very first replication of the new genotype, and the smaller its selective advantage, the more likely it is not to replicate even once. Now, if a new mutant with a poor replication rate w_{new} but high effective fidelity Q_{new} arises in a population that is dominated by sequences with large intrinsic replication rate, we would intuitively assume that the mutant will hardly ever replicate even once, and therefore will never get a chance to employ its superior fidelity. However, this is not correct if the effective fidelity of the dominating sequences, Q_{dom} , is low. From Eq. (4), we find that the concentration of sequences that actually replicate is given by Q_{dom} . Therefore, even though the sequences that replicate do so at a high rate, the actual number of births that occur is small, comparable to the one in a population in which all individuals reproduce with rate $w_{\text{dom}}Q_{\text{dom}}$. Therefore, the newly introduced genotype is relatively safe from being washed out prematurely, and fixation takes place at the predicted rate.

CONCLUSIONS

We have demonstrated that for a population in a landscape where neutral mutants abound, the product of intrinsic replication rate w and effective copy-fidelity Q is being maximized under selection, rather than the intrinsic replication rate alone. This observation has led to the natural distinction between two modes of selection, one in which intrinsic replication rate is favored, and one in which robustness (high Q) is more important. In the latter phase, the success of a single sequence depends strongly on the mutant cloud the sequence belongs to. Our results thus demonstrate that the unit of selection in molecular evolution is indeed the quasispecies, as proposed by Eigen and Schuster (1979), and not the individual replicating sequence. In particular, the probability of

fixation of a single advantageous mutant in an established quasispecies can be predicted accurately with results from standard population genetics, provided we consider the overall growth rates of the established quasispecies and the quasispecies potentially formed by the mutant, rather than the replication rates of mutant and established wild type.

Acknowledgments

This work was supported by the NSF under contract No DEB-9981397. C.O.W. would like to thank (in alphabetical order) C. Adami for many useful comments and suggestions; P. Campos for double-checking fixation probabilities; W. Fontana for providing the original flow-reactor code; J. Wang for writing an early Mathematica script used in this study.

References

- Alves, D., P. R. A. Campos, A. T. C. Silva, and J. F. Fontanari. 2001. Group selection models in prebiotic evolution. *Phys. Rev. E* 63, 011911.
- Ancel, L. W. and W. Fontana 2000. Plasticity, evolvability, and modularity in RNA. *J. of Exp. Zoology* 288, 242–283.
- Biebricher, C. K. 1987. Replication and evolution of short-chained RNA species replicated by Q β replicase. *Cold Spring Harbor Symp. Quant. Biol.* 52, 299–306.
- Biebricher, C. K. and W. C. Gardiner. 1997. Molecular evolution of RNA in vitro. *Biophys. Chem.* 66, 179–192.
- Burch, C. L. and L. Chao. 2000. Evolvability of an RNA virus is determined by its mutational neighbourhood. *Nature* 406, 625–628.
- Domingo, E. and J. J. Holland. 1997. RNA virus mutations and fitness for survival. *Annu. Rev. Microbiol.* 51, 151–178.
- Drake, J. W. 1993. Rates of spontaneous mutation among RNA viruses. *Proc. Natl. Acad. Sci. USA* 90, 4171–4175.
- Eigen, M., J. McCaskill, and P. Schuster. 1988. Molecular quasi-species. *J. Phys. Chem.* 92, 6881–6891.
- . 1989. The molecular quasi-species. *Adv. Chem. Phys.* 75, 149–263.
- Eigen, M. and P. Schuster. 1979. *The Hypercycle—A Principle of Natural Self-Organization*. Springer, Berlin.
- Ewens, W. J. 1979. *Mathematical Population Genetics*. Springer, Berlin.
- Feller, W. 1968. *An Introduction to Probability Theory and Its Applications* (3rd ed.), Volume I. Wiley, New York.
- Fontana, W. and P. Schuster. 1998. Continuity in evolution: on the nature of transitions. *Science* 280, 1451–1455.
- Fontana, W., P. F. Stadler, E. G. Bornberg-Bauer, T. Griesmacher, I. L. Hofacker, M. Tacker, P. Tarazona, E. D. Weinberger, and P. Schuster. 1993. RNA folding and combinatorial landscapes. *Phys. Rev. E* 47, 2083–2099.
- Gavrilets, S. 1997. Evolution and speciation on holey adaptive landscapes. *TREE* 12, 307–312.
- Gavrilets, S. and J. Gravner. 1997. Percolation on the fitness hypercube and the evolution of reproductive isolation. *J. theor. Biol.* 184, 51–64.
- Gillespie, D. T. 1976. A general method for numerically simulating the stochastic time evolution of coupled chemical reactions. *J. Comput. Phys.* 22, 403–434.
- Haldane, J. B. S. 1927. A mathematical theory of natural and artificial selection. Part V: Selection and mutation. *Proc. Camb. Phil. Soc.* 23, 838–844.
- Hofacker, I. L., W. Fontana, P. F. Stadler, S. Bonhoeffer, M. Tacker, and P. Schuster. 1994. Fast folding and comparison of RNA secondary structures. *Monatshefte f. Chemie* 125, 167–188.
- Huynen, M. A., P. F. Stadler, and W. Fontana. 1996. Smoothness within ruggedness: The role of neutrality in adaptation. *Proc. Natl. Acad. Sci. USA* 93, 397–401.
- Kimura, M. 1964. Diffusion models in population genetics. *J. Appl. Prob.* 1, 177–232.
- Lenski, R. E. and B. R. Levin. 1985. Constraints on the coevolution of bacteria and virulent phage:

- a model, some experiments, and predictions for natural communities. *Am. Nat.* 125, 585–602.
- Maynard Smith, J. 1993. *The Theory of Evolution* (3rd ed.). Cambridge University Press, New York.
- Moya, A., S. F. Elena, A. Bracho, R. Miralles, and E. Barrio. 2000. The evolution of RNA viruses: A population genetics view. *Proc. Natl. Acad. Sci. USA* 97, 6967–6973.
- Nowak, M. A. 1992. What is a quasispecies? *TREE* 7, 118–121.
- Ofria, C. and C. Adami. 2001. Evolution of genetic organization in digital organisms. In L. Landweber and E. Winfree (Eds.), *Evolution as Computation*, pp. 167. Springer, New York.
- Reidys, C., C. V. Forst, and P. Schuster. 2001. Replication and mutation on neutral networks. *Bull. Math. Biol.* 63, 57–94.
- Schuster, P. and W. Fontana. 1999. Chance and necessity in evolution: lessons from RNA. *Physica D* 133, 427–452.
- Schuster, P. and J. Swetina. 1988. Stationary mutant distributions and evolutionary optimization. *Bull. Math. Biol.* 50, 635–660.
- Stanley, E. H. 1971. *Introduction to Phase Transitions and Critical Phenomena*. Oxford University Press, New York.
- Steinhauer, D. A., J. C. de la Torre, E. Meier, and J. J. Holland. 1989. Extreme heterogeneity in populations of vesicular stomatitis virus. *J. of Virology* 63, 2072–2080.
- van Nimwegen, E., J. P. Crutchfield, and M. Huyenen. 1999. Neutral evolution of mutational robustness. *Proc. Natl. Acad. Sci. USA* 96, 9716–9720.
- Walter, A. E., D. H. Turner, J. Kim, M. H. Lyttle, P. Müller, D. H. Mathews, and M. Zuker. 1994. Coaxial stacking of helices enhances binding of oligoribonucleotides and improves predictions of RNA folding. *Proc. Natl. Acad. Sci. USA* 91, 9218–9222.
- Wilke, C. O., C. Ronnewinkel, and T. Martinetz. 2001. Dynamic fitness landscapes in molecular evolution. *Phys. Rep.* in press.
- Wilson, D. S. 1979. *The Natural Selection of Populations & Communities*. Benjamin-Cummings, Menlo Park.

Figure 1: The two different folds used in this study. Both consist of the same number of base pairs ($l = 62$), but Fold 1 has a higher neutrality ($\nu = 0.442$) than Fold 2 ($\nu = 0.366$). See also Fig. 2.

Figure 2: Decay of the steady state concentration x_1 as a function of $1 - q$ for two example secondary structures. The solid and the dashed line are given by $\exp[-l(1 - q)(1 - \nu_i)]$ with $l = 62$. The values for ν_1 and ν_2 have been obtained from a fit of this expression to the measured data (shown as points with bars indicating the standard error).

Figure 3: Typical phase diagram following from Eq. 11. We used $l = 100$, $w_2 = 1$, and $\nu_1 = 0.5$, as well as $\nu_2 = 0.6$ in graph a) and $w_1 = 1.5$ in graph b).

Figure 4: Concentrations $x_1(t)$ and $x_2(t)$ as functions of the time t for a copy fidelity of $q = 0.99$. The thick lines represent the analytic predictions from Eqs. (10a) and (10b), and the thin lines stem from simulations with $N = 1000$.

Figure 5: Concentrations $x_1(200)$ (dashed lines) and $x_2(200)$ (solid lines) as functions of the per-nucleotide mutation rate $1 - q$. The lines represent the analytic predictions. The points represent the average over 25 independent simulation runs each, with bars indicating the standard error. We performed the simulations with four different population sizes, $N = 5000$ (a), $N = 1000$ (b), $N = 500$ (c), and $N = 100$ (d). The initial concentrations in all simulations were $x_1(0) = x_2(0) = 0.5$, $x_d(0) = 0$.

Figure 6: Probability of fixation as a function of the mutation rate. Below $1 - q = 0.02$, we are looking at the probability of fixation of a single sequence of type 2 in a full population of sequences of type 1. Above $1 - q = 0.02$, we are considering the reversed configuration. The solid and dashed line represent the analytical prediction $\pi = s/(1 + s)$, the points stem from simulations (bars indicate standard error).

Figure 1:






Figure 2:







Figure 3:





Figure 4:

Figure 5:






Figure 6: